



Invasive Species Management Plans for Florida

Air Potato

Dioscorea bulbifera (L.) Dioscoreaceae

INTRODUCTION

A native to tropical Asia, air potato, *Dioscorea bulbifera*, was first introduced to the Americas from Africa. In 1905 it was introduced to Florida. Due to its ability to displace native species and disrupt natural processes such as fire and water flow, air potato has been listed as one of Florida's most invasive plant species since 1993, and was placed on the Florida Noxious Weed List by the Florida Department of Agriculture and Consumer Services in 1999.

DESCRIPTION

Air potato is in the family Dioscoreaceae, or simply the Yam Family. It is an herbaceous twining vine, growing 70 feet or more in length. Leaves are broadly cordate (heart shaped) and alternately arranged on stems. A distinguishing characteristic of air potato is that all leaf veins arise from the leaf base, unlike other herbaceous vines such as smilax and morningglories. Flowers are inconspicuous, arising from leaf axils in panicles 4 inches long, and are fairly uncommon in Florida. Vegetative reproduction is the primary mechanism of spread. This is through the formation of aerial tubers, or bulbils, which are formed in leaf axils. These vary in roundish shapes and sizes. In addition, large tubers are formed underground, some reaching over 6 inches in diameter.

Dioscorea alata or winged yam can easily be mistaken for air potato, *D. bulbifera*. Winged yam gets its name from its winged internodes, a characteristic feature of the species. Another difference between *D. alata* and *D. bulbifera* is the arrangement of the leaves. *D. alata* has opposite leaves as opposed to the alternate leaves of air potato. Winged yam grows to 30 feet, roughly half the length of the invasive species. This species of *Dioscorea* does not produce nearly as many bulbils as *D. bulbifera*. However, this species can also be considered invasive and problematic, but to a lesser extent than *D. bulbifera*. Although considered to be a species of yam, these plants are very toxic and **should not be consumed.**

IMPACTS

Air potato can grow extremely quickly, roughly 8 inches per day. It typically climbs to the tops of trees and has a tendency to take over native plants. New plants develop from bulbils that form on the plant, and these bulbils serve as a means of dispersal. The aerial stems of air potato die back in winter, but resprouting occurs from bulbils and underground tubers. The primary means of spread and reproduction are via bulbils. The smallest bulbils make control of air potato difficult due to their ability to sprout at a very small stage.

MANAGEMENT

Preventative: Prevention is a key step in the management of air potato. Bulbils are the primary mechanism of spread, and research has shown even minutely small propagules can sprout and form new plants. How these bulbils are spread is speculative, but it appears movement of contaminated brush, debris or soil is the primary mechanism. Mowers and other brush-cutting equipment may also disperse long distances, either through contaminated equipment or throwing of the bulbils during the mowing operation. Spread via birds and other animals may occur, but this has not been confirmed. Water is also a major means of dispersal, so care must be taken to first eliminate populations along water bodies where bulbils may be easily spread. In addition, extra time must be utilized after flood events, as spread may be extensive.

Cultural: Weeds such as air potato generally invade open or disturbed areas – following a burn, clearing mowing, etc., so these areas are particularly vulnerable to invasion. Therefore, a healthy ecosystem with good species diversity will help to deter infestation.

Another very important combined cultural and mechanical method is the air potato roundup. Each year many counties in Florida, including Hernando, Alachua, and Duval counties, recruit volunteers to help protect and conserve Florida's natural areas through the removal of air potato. During the air potato round up, citizens, organizations, and local businesses get together to collect vines and bulbils. In 2003, the City of Gainesville collected 13 tons of air potato and other invasive plants (Gainesville Parks and Recreation). Removing bulbils and vines from natural areas helps prevent the spread of

air potato to new areas, as well as reduces the possibility of reinfestation. In addition to collecting and removing aboveground bulbils, digging up and removing below ground tubers will help. This may be particularly useful to eliminate isolated plants/small populations – especially in areas that cannot be easily accessed or chemically treated. One of the most important control measures for air potato is the removal of bulbils and tubers.

Mechanical: Mechanical methods are limited for air potato, as control of the vines generally results in damage to the vegetation being climbed/smothered by the air potato. Burning also results in excessive damage to the native vegetation, as the fire follows the vines into the tree canopy. Mowing will help to suppress air potato, but as mentioned previously, this may increase the overall problem due to spreading of the bulbils.

Biological: There is limited research and data on biological control of air potato.

Chemical: Chemical control is one of the most effective means of control for air potato, but single applications will generally not provide complete control. This is due to resprouting of bulbils or underground tubers. A dilution of triclopyr (Garlon 3A at 1 to 2% solution or Garlon 4 at 0.5 to 2% solution) in water can be an effective control for air potato when applied as a foliar application. Be sure to include a non-ionic surfactant at 0.25% (10 mLs or 2 teaspoons per gallon of spray solution). A 2 to 3% solution of glyphosate (Roundup, etc.) can also be effective. These herbicides are systemic (move throughout plant tissue) so care must be exercised to minimize off-target damage. If air potato vines are growing up into trees or other desirable species, vines should be cut or pulled down to minimize damage to the desirable vegetation. Pulling the vines down without severing them from the underground tuber will allow the herbicide to move into the tuber and provide better control. The best time to apply an herbicide is in the spring and summer when air potato is actively growing. Be sure to allow adequate time for the plant to regrow from the winter to ensure movement of the herbicide back into the underground tuber. (As plants grow and mature, they begin to move sugars back into the roots and below-ground tubers). However, treat *before* the plants begin to form new bulbils. Persistence and integration of control methods will be the key to complete air potato management.

REFERENCES:

Langeland, K.A. 2003. Natural Area Weeds: Air Potato (*Dioscorea bulbifera*). IFAS Publication SS AGR 164. Florida Cooperative Extension Service, Agronomy Department, University of Florida.

Langeland, K.A. and K. Craddock Burks. 1998. Identification and Biology of Non-Native Plants in Florida's Natural Areas. IFAS Publication SP 257. University of Florida, Gainesville. 165 pp.

Langeland K.A. and R.K. Stocker. 2001. Control of Non-Native Plants in Natural Areas of Florida. IFAS Publication SP 242. University of Florida, Gainesville. 34pp.

HELPFUL LINKS:

Center for Aquatic and Invasive Plants Web Site: <http://www.plants.ifas.ufl.edu>

Nature Operations Division, Gainesville Parks and Recreation:
<http://www.natureoperations.org/>

Florida Exotic Pest Plant Council: <http://www.fleppc.org/>

Mature Plant

- Rapidly climbing, twining herbaceous vine
- Vines killed by frost
- Regrowth from underground tubers



Leaves and Flowers

- Leaves cordate
- All leaf veins arise from leaf base
- Flowers are inconspicuous, arise from panicles from leaf axils



Bulbils

- Aerial tubers, borne in leaf axils
- Generally roundish, smooth and gray to brown in color
- Primary means of spread

